

1.5 Functions and Logarithms

What you will learn about . . .

- One-to-One Functions
- Inverses
- Finding Inverses
- Logarithmic Functions
- Properties of Logarithms
- Applications

and why . . .

Logarithmic functions are used in many applications, including finding time in investment problems.

One-to-One Functions

As you know, a function is a rule that assigns a single value in its range to each point in its domain. Some functions assign the same output to more than one input. For example, $f(x) = x^2$ assigns the output 4 to both 2 and -2. Other functions never output a given value more than once. For example, the cubes of different numbers are always different.

If each output value of a function is associated with exactly one input value, the function is *one-to-one*.

DEFINITION One-to-One Function

A function $f(x)$ is **one-to-one** on a domain D if $f(a) \neq f(b)$ whenever $a \neq b$.

The graph of a one-to-one function $y = f(x)$ can intersect any horizontal line at most once (the *horizontal line test*). If it intersects such a line more than once it assumes the same y -value more than once, and is therefore not one-to-one (Figure 1.33).




Figure 1.33 Using the horizontal line test, we see that $y = x^3$ is one-to-one and $y = x^2$ is not.

EXAMPLE 1 Using the Horizontal Line Test

Determine whether the functions are one-to-one.

- (a) $f(x) = |x|$ (b) $g(x) = \sqrt{x}$

SOLUTION

(a) As Figure 1.34a suggests, each horizontal line $y = c$, $c > 0$, intersects the graph of $f(x) = |x|$ twice. So f is not one-to-one.

(b) As Figure 1.34b suggests, each horizontal line intersects the graph of $g(x) = \sqrt{x}$ either once or not at all. The function g is one-to-one.

Now Try Exercise 1.




Figure 1.34 (a) The graph of $f(x) = |x|$ and a horizontal line. (b) The graph of $g(x) = \sqrt{x}$ and a horizontal line. (Example 1)

Inverses

Since each output of a one-to-one function comes from just one input, a one-to-one function can be reversed to send outputs back to the inputs from which they came. The function

defined by reversing a one-to-one function f is the **inverse of f** . The functions in Tables 1.13 and 1.14 are inverses of one another. The symbol for the inverse of f is f^{-1} , read "f inverse." The -1 in f^{-1} is not an exponent; $f^{-1}(x)$ does not mean $1/f(x)$.

TABLE 1.13 Rental Charge versus Time	
Time x (hours)	Charge y (dollars)
1	5.00
2	7.50
3	10.00
4	12.50
5	15.00
6	17.50

TABLE 1.14 Time versus Rental Charge	
Charge x (dollars)	Time y (hours)
5.00	1
7.50	2
10.00	3
12.50	4
15.00	5
17.50	6

As Tables 1.13 and 1.14 suggest, composing a function with its inverse in either order sends each output back to the input from which it came. In other words, the result of composing a function and its inverse in either order is the **identity function**, the function that assigns each number to itself. This gives a way to test whether two functions f and g are inverses of one another. Compute $f \circ g$ and $g \circ f$. If $(f \circ g)(x) = (g \circ f)(x) = x$, then f and g are inverses of one another; otherwise they are not. The functions $f(x) = x^3$ and $g(x) = x^{1/3}$ are inverses of one another because $(x^3)^{1/3} = x$ and $(x^{1/3})^3 = x$ for every number x .

EXPLORATION 1 Testing for Inverses Graphically

For each of the function pairs below,

(a) Graph f and g together in a square window.

(b) Graph $f \circ g$. (c) Graph $g \circ f$.


What can you conclude from the graphs?

1. $f(x) = x^3$, $g(x) = x^{1/3}$
2. $f(x) = x$, $g(x) = 1/x$
3. $f(x) = 3x$, $g(x) = x/3$
4. $f(x) = e^x$, $g(x) = \ln x$

Finding Inverses

How do we find the graph of the inverse of a function? Suppose, for example, that the function is the one pictured in Figure 1.35a. To read the graph, we start at the point x on the x -axis, go up to the graph, and then move over to the y -axis to read the value of y . If we start with y and want to find the x from which it came, we reverse the process (Figure 1.35b).

The graph of f is already the graph of f^{-1} , although the latter graph is not drawn in the usual way with the domain axis horizontal and the range axis vertical. For f^{-1} , the input-output pairs are reversed. To display the graph of f^{-1} in the usual way, we have to reverse the pairs by reflecting the graph across the 45° line $y = x$ (Figure 1.35c) and interchanging the letters x and y (Figure 1.35d). This puts the independent variable, now called x , on the horizontal axis and the dependent variable, now called y , on the vertical axis.

Figure 1.35 The graph of $y = f^{-1}(x)$.

The fact that the graphs of f and f^{-1} are reflections of each other across the line $y = x$ is to be expected because the input-output pairs (a, b) of f have been reversed to produce the input-output pairs (b, a) of f^{-1} .

The pictures in Figure 1.35 tell us how to express f^{-1} as a function of x algebraically.

Writing f^{-1} as a Function of x

1. Solve the equation $y = f(x)$ for x in terms of y .
2. Interchange x and y . The resulting formula will be $y = f^{-1}(x)$.

EXAMPLE 2 Finding the Inverse Function

Show that the function $y = f(x) = -2x + 4$ is one-to-one and find its inverse function.

SOLUTION

Every horizontal line intersects the graph of f exactly once, so f is one-to-one and has an inverse.

Step 1:

Solve for x in terms of y : $y = -2x + 4$

$$x = -\frac{1}{2}y + 2$$

continued

Step 2:

Interchange x and y : $y = -\frac{1}{2}x + 2$

The inverse of the function $f(x) = -2x + 4$ is the function $f^{-1}(x) = -(1/2)x + 2$. We can verify that both composites are the identity function.

$$f^{-1}(f(x)) = -\frac{1}{2}(-2x + 4) + 2 = x - 2 + 2 = x$$

$$f(f^{-1}(x)) = -2\left(-\frac{1}{2}x + 2\right) + 4 = x - 4 + 4 = x$$

Now Try Exercise 13.


Graphing $y = f(x)$ and $y = f^{-1}(x)$ Parametrically

We can graph any function $y = f(x)$ as

$$x_1 = t, \quad y_1 = f(t).$$

Interchanging t and $f(t)$ produces parametric equations for the inverse:

$$x_2 = f(t), \quad y_2 = t.$$



$[-1.5, 3.5]$ by $[-1, 2]$

Figure 1.36 The graphs of f and f^{-1} are reflections of each other across the line $y = x$. (Example 3)

We can use parametric graphing to graph the inverse of a function without finding an explicit rule for the inverse, as illustrated in Example 3.

EXAMPLE 3 Graphing the Inverse Parametrically

(a) Graph the one-to-one function $f(x) = x^2$, $x \geq 0$, together with its inverse and the line $y = x$, $x \geq 0$.

(b) Express the inverse of f as a function of x .

SOLUTION

(a) We can graph the three functions parametrically as follows:

Graph of f : $x_1 = t, \quad y_1 = t^2, \quad t \geq 0$

Graph of f^{-1} : $x_2 = t^2, \quad y_2 = t$

Graph of $y = x$: $x_3 = t, \quad y_3 = t$

Figure 1.36 shows the three graphs.

(b) Next we find a formula for $f^{-1}(x)$.

Step 1:

Solve for x in terms of y .

$$\begin{aligned} y &= x^2 \\ \sqrt{y} &= \sqrt{x^2} \\ \sqrt{y} &= x && \text{Because } x \geq 0 \end{aligned}$$

Step 2:

Interchange x and y .

$$\sqrt{x} = y$$

Thus, $f^{-1}(x) = \sqrt{x}$.

Now Try Exercise 27.


Logarithmic Functions

If a is any positive real number other than 1, the base a exponential function $f(x) = a^x$ is one-to-one. It therefore has an inverse. Its inverse is called the *base a logarithm function*.

DEFINITION Base a Logarithm Function

The **base a logarithm function** $y = \log_a x$ is the inverse of the base a exponential function $y = a^x$ ($a > 0, a \neq 1$).

The domain of $\log_a x$ is $(0, \infty)$, the range of a^x . The range of $\log_a x$ is $(-\infty, \infty)$, the domain of a^x .



[-6, 6] by [-4, 4]

Figure 1.37 The graphs of $y = 2^x$ ($x_1 = t$, $y_1 = 2^t$), its inverse $y = \log_2 x$ ($x_2 = 2^t$, $y_2 = t$), and $y = x$ ($x_3 = t$, $y_3 = t$).

Because we have no technique for solving for x in terms of y in the equation $y = a^x$, we do not have an explicit formula for the logarithm function as a function of x . However, the graph of $y = \log_a x$ can be obtained by reflecting the graph of $y = a^x$ across the line $y = x$, or by using parametric graphing (Figure 1.37).

Logarithms with base e and base 10 are so important in applications that calculators have special keys for them. They also have their own special notation and names:

$$\log_e x = \ln x,$$

$$\log_{10} x = \log x$$

The function $y = \ln x$ is called the **natural logarithm function** and $y = \log x$ is often called the **common logarithm function**.

Properties of Logarithms

Because a^x and $\log_a x$ are inverses of each other, composing them in either order gives the identity function. This gives two useful properties.

Inverse Properties for a^x and $\log_a x$

1. Base a : $a^{\log_a x} = x$, $\log_a a^x = x$, $a > 1$, $x > 0$
2. Base e : $e^{\ln x} = x$, $\ln e^x = x$, $x > 0$

These properties help us with the solution of equations that contain logarithms and exponential functions.

EXAMPLE 4 Using the Inverse Properties

Solve for x : (a) $\ln x = 3t + 5$ (b) $e^{2x} = 10$

SOLUTION

(a) $\ln x = 3t + 5$

$$e^{\ln x} = e^{3t+5} \quad \text{Exponentiate both sides.}$$

$$x = e^{3t+5} \quad \text{Inverse Property}$$

(b) $e^{2x} = 10$

$$\ln e^{2x} = \ln 10 \quad \text{Take logarithms of both sides.}$$

$$2x = \ln 10 \quad \text{Inverse Property}$$

$$x = \frac{1}{2} \ln 10 \approx 1.15$$

Now Try Exercises 33 and 37.

The logarithm function has the following useful arithmetic properties.

Properties of Logarithms

For any real numbers $x > 0$ and $y > 0$,

1. **Product Rule:** $\log_a xy = \log_a x + \log_a y$
2. **Quotient Rule:** $\log_a \frac{x}{y} = \log_a x - \log_a y$
3. **Power Rule:** $\log_a x^y = y \log_a x$

EXPLORATION 2 Supporting the Product Rule

Let $y_1 = \ln(ax)$, $y_2 = \ln x$, and $y_3 = y_1 - y_2$.


1. Graph y_1 and y_2 for $a = 2, 3, 4$, and 5. How do the graphs of y_1 and y_2 appear to be related?
2. Support your finding by graphing y_3 .
3. Confirm your finding algebraically.

The following formula allows us to evaluate $\log_a x$ for any base $a > 0$, $a \neq 1$, and to obtain its graph using the natural logarithm function on our grapher.

Change of Base Formula

$$\log_a x = \frac{\ln x}{\ln a}$$

$$y = \frac{\ln x}{\ln 2}$$



[-6, 6] by [-4, 4]

Figure 1.38 The graph of $f(x) = \log_2 x$ using $f(x) = (\ln x)/(\ln 2)$. (Example 5)

EXAMPLE 5 Graphing a Base a Logarithm Function

Graph $f(x) = \log_2 x$.

SOLUTION

We use the change of base formula to rewrite $f(x)$.

$$f(x) = \log_2 x = \frac{\ln x}{\ln 2}$$

Figure 1.38 gives the graph of f .

Now Try Exercise 41.

Applications

In Section 1.3 we used graphical methods to solve exponential growth and decay problems. Now we can use the properties of logarithms to solve the same problems algebraically.

EXAMPLE 6 Finding Time

Sarah invests \$1000 in an account that earns 5.25% interest compounded annually. How long will it take the account to reach \$2500?

SOLUTION

Model The amount in the account at any time t in years is $1000(1.0525)^t$, so we need to solve the equation

$$1000(1.0525)^t = 2500.$$


continued

TABLE 1.15 Saudi Arabia's Natural Gas Production

Year	Cubic Feet (trillions)
2002	2.00
2003	2.12
2004	2.32
2005	2.52
2006	2.59

Source: Statistical Abstract of the United States, 2010.

$$f(x) = 1.558 + (0.571) \ln x$$



[-2, 10] by [-2, 4]

Figure 1.39 The value of f at $x = 8$ is about 2.75. (Example 7)

Solve Algebraically

$$(1.0525)^t = 2.5$$

$$\ln (1.0525)^t = \ln 2.5$$

$$t \ln 1.0525 = \ln 2.5$$

$$t = \frac{\ln 2.5}{\ln 1.0525} \approx 17.9$$

Divide by 1000.

Take logarithms of both sides.

Power Rule

Interpret The amount in Sarah's account will be \$2500 in about 17.9 years, or about 17 years and 11 months.

Now Try Exercise 47.

EXAMPLE 7 Estimating Natural Gas Production

Table 1.15 shows the annual number of cubic feet in trillions of natural gas produced by Saudi Arabia for several years.

Find the natural logarithm regression equation for the data in Table 1.15 and use it to estimate the number of cubic feet of natural gas produced by Saudi Arabia in 2008.

SOLUTION

Model We let $x = 0$ represent 2000, $x = 1$ represent 2001, and so forth. We compute the natural logarithm regression equation to be

$$f(x) = 1.558 + 0.571 \ln(x).$$

Solve Graphically Figure 1.39 shows the graph of f superimposed on the scatter plot of the data. The year 2008 is represented by $x = 8$. Reading from the graph we find $f(8) = 2.75$ trillion cubic feet.

Now Try Exercise 49.

Quick Review 1.5 (For help, go to Sections 1.2, 1.3, and 1.4.)

Exercise numbers with a gray background indicate problems that authors have designed to be solved *without a calculator*.

In Exercises 1–4, let $f(x) = \sqrt[3]{x-1}$, $g(x) = x^2 + 1$, and evaluate the expression.

1. $(f \circ g)(1)$
2. $(g \circ f)(-7)$
3. $(f \circ g)(x)$
4. $(g \circ f)(x)$

In Exercises 5 and 6, choose parametric equations and a parameter interval to represent the function on the interval specified.


5. $y = \frac{1}{x-1}$, $x \geq 2$
6. $y = x$, $x < -3$

In Exercises 7–10, find the points of intersection of the two curves. Round your answers to 2 decimal places.

7. $y = 2x - 3$, $y = 5$
8. $y = -3x + 5$, $y = -3$
9. (a) $y = 2^x$, $y = 3$
(b) $y = 2^x$, $y = -1$
10. (a) $y = e^{-x}$, $y = 4$
(b) $y = e^{-x}$, $y = -1$

Section 1.5 Exercises

In Exercises 1–6, determine whether the function is one-to-one.



In Exercises 7–12, determine whether the function has an inverse function.

7. $y = \frac{3}{x-2} - 1$
8. $y = x^2 + 5x$
9. $y = x^3 - 4x + 6$
10. $y = x^3 + x$
11. $y = \ln x^2$
12. $y = 2^{3-x}$

In Exercises 13–24, find f^{-1} and verify that

$$(f \circ f^{-1})(x) = (f^{-1} \circ f)(x) = x.$$

13. $f(x) = 2x + 3$
14. $f(x) = 5 - 4x$
15. $f(x) = x^3 - 1$
16. $f(x) = x^2 + 1$, $x \geq 0$
17. $f(x) = x^2$, $x \leq 0$
18. $f(x) = x^{2/3}$, $x \geq 0$
19. $f(x) = -(x-2)^2$, $x \leq 2$
20. $f(x) = x^2 + 2x + 1$, $x \geq -1$
21. $f(x) = \frac{1}{x^2} + 1$, $x > 0$
22. $f(x) = \frac{1}{x^3} + 3$
23. $f(x) = \frac{2x+1}{x+3}$
24. $f(x) = \frac{x+3}{x-2}$

In Exercises 25–32, use parametric graphing to graph f , f^{-1} , and $y = x$.

25. $f(x) = e^x$
26. $f(x) = 3^x$
27. $f(x) = 2^{-x}$
28. $f(x) = 3^{-x}$
29. $f(x) = \ln x$
30. $f(x) = \log x$
31. $f(x) = \sin^{-1} x$
32. $f(x) = \tan^{-1} x$

In Exercises 33–36, solve the equation algebraically. Support your solution graphically.

33. $(1.045)^t = 2$
34. $e^{0.05t} = 3$
35. $e^x + e^{-x} = 3$
36. $2^x + 2^{-x} = 5$

In Exercises 37 and 38, solve for y .

37. $\ln y = 2t + 4$
38. $\ln(y-1) - \ln 2 = x + \ln x$

In Exercises 39–42, draw the graph and determine the domain and range of the function.

39. $y = 2 \ln(3-x) - 4$
40. $y = -3 \log(x+2) + 1$
41. $y = \log_2(x+1)$
42. $y = \log_3(x-4)$

In Exercises 43 and 44, find a formula for f^{-1} and verify that $(f \circ f^{-1})(x) = (f^{-1} \circ f)(x) = x$.

43. $f(x) = \frac{100}{1+2^{-x}}$
44. $f(x) = \frac{50}{1+1.1^{-x}}$

45. **Self-inverse** Prove that the function f is its own inverse.

- (a) $f(x) = \sqrt{1-x^2}$, $x \geq 0$
- (b) $f(x) = 1/x$

46. **Radioactive Decay** The half-life of a certain radioactive substance is 12 hours. There are 8 grams present initially.

- (a) Express the amount of substance remaining as a function of time t .
- (b) When will there be 1 gram remaining?

47. **Doubling Your Money** Determine how much time is required for a \$500 investment to double in value if interest is earned at the rate of 4.75% compounded annually.

48. **Population Growth** The population of Glenbrook is 375,000 and is increasing at the rate of 2.25% per year. Predict when the population will be 1 million.

In Exercises 49 and 50, let $x = 0$ represent 1990, $x = 1$ represent 1991, and so forth.

49. Natural Gas Production

- (a) Find a natural logarithm regression equation for the data in Table 1.16 and superimpose its graph on a scatter plot of the data. Let $x = 0$ represent 2000.

TABLE 1.16 Iran's Natural Gas Production

Year	Cubic Feet (trillions)
2002	2.65
2003	2.86
2004	2.96
2005	3.56
2006	3.84

Source: Statistical Abstract of the United States, 2010.

- (b) Estimate the number of cubic feet of natural gas produced by Iran in 2008.
- (c) Using the model in part (b), predict when Iran's natural gas production reaches 4.2 trillion cubic feet.

50. Natural Gas Production

- (a) Find a natural logarithm regression equation for the data in Table 1.17 and superimpose its graph on a scatter plot of the data. Let $x = 0$ represent 2000.

TABLE 1.17
China's Natural Gas Production


Year	Cubic Feet (trillions)
2002	1.15
2003	1.21
2004	1.44
2005	1.76
2006	2.07

Source: Statistical Abstract of the United States, 2010.

- (b) Estimate the number of cubic feet of natural gas produced by China in 2008.
- (c) Using the model in part (b), predict when China's natural gas production reaches 2.25 trillion cubic feet.
- 51. Group Activity Inverse Functions** Let $y = f(x) = mx + b$, $m \neq 0$.
- (a) **Writing to Learn** Give a convincing argument that f is a one-to-one function.
- (b) Find a formula for the inverse of f . How are the slopes of f and f^{-1} related?
- (c) If the graphs of two functions are parallel lines with a nonzero slope, what can you say about the graphs of the inverses of the functions?
- (d) If the graphs of two functions are perpendicular lines with a nonzero slope, what can you say about the graphs of the inverses of the functions?

Standardized Test Questions

- 52. True or False** The function displayed in the graph below is one-to-one. Justify your answer.



- 53. True or False** If $(f \circ g)(x) = x$, then g is the inverse function of f . Justify your answer.

In Exercises 54 and 55, use the function $f(x) = 3 - \ln(x + 2)$.

- 54. Multiple Choice** Which of the following is the domain of f ?

(A) $x \neq -2$ (B) $(-\infty, \infty)$ (C) $(-2, \infty)$
(D) $[-1.9, \infty)$ (E) $(0, \infty)$

- 55. Multiple Choice** Which of the following is the range of f ?

(A) $(-\infty, \infty)$ (B) $(-\infty, 0)$ (C) $(-2, \infty)$
(D) $(0, \infty)$ (E) $(0, 5.3)$

- 56. Multiple Choice** Which of the following is the inverse of $f(x) = 3x - 2$?

(A) $g(x) = \frac{1}{3x-2}$ (B) $g(x) = x$ (C) $g(x) = 3x - 2$
(D) $g(x) = \frac{x-2}{3}$ (E) $g(x) = \frac{x+2}{3}$

- 57. Multiple Choice** Which of the following is a solution of the equation $2 - 3^{-x} = -1$?

(A) $x = -2$ (B) $x = -1$ (C) $x = 0$
(D) $x = 1$ (E) There are no solutions.

Exploration

- 58. Supporting the Quotient Rule** Let $y_1 = \ln(x/a)$, $y_2 = \ln x$, $y_3 = y_2 - y_1$, and $y_4 = e^{y_3}$.

- (a) Graph y_1 and y_2 for $a = 2, 3, 4$, and 5 . How are the graphs of y_1 and y_2 related?
- (b) Graph y_3 for $a = 2, 3, 4$, and 5 . Describe the graphs.
- (c) Graph y_4 for $a = 2, 3, 4$, and 5 . Compare the graphs to the graph of $y = a$.
- (d) Use $e^{y_3} = e^{y_2 - y_1} = a$ to solve for y_1 .

Extending the Ideas

- 59. One-to-One Functions** If f is a one-to-one function, prove that $g(x) = -f(x)$ is also one-to-one.
- 60. One-to-One Functions** If f is a one-to-one function and $f(x)$ is never zero, prove that $g(x) = 1/f(x)$ is also one-to-one.
- 61. Domain and Range** Suppose that $a \neq 0$, $b \neq 1$, and $b > 0$. Determine the domain and range of the function.
- (a) $y = a(b^{c-x}) + d$ (b) $y = a \log_b(x - c) + d$

62. Group Activity Inverse Functions

Let $f(x) = \frac{ax+b}{cx+d}$, $c \neq 0$, $ad - bc \neq 0$.

- (a) **Writing to Learn** Give a convincing argument that f is one-to-one.
- (b) Find a formula for the inverse of f .
- (c) Find the horizontal and vertical asymptotes of f .
- (d) Find the horizontal and vertical asymptotes of f^{-1} . How are they related to those of f ?

1.6 Trigonometric Functions**Radian Measure**

The **radian measure** of the angle ACB at the center of the unit circle (Figure 1.40) equals the length of the arc that ACB cuts from the unit circle.

EXAMPLE 1 Finding Arc Length

Find the length of an arc subtended on a circle of radius 3 by a central angle of measure $2\pi/3$.

SOLUTION

According to Figure 1.40, if s is the length of the arc, then

$$s = r\theta = 3(2\pi/3) = 2\pi.$$

Now Try Exercise 1.

What you will learn about . . .

- Radian Measure
- Graphs of Trigonometric Functions
- Periodicity
- Even and Odd Trigonometric Functions
- Transformations of Trigonometric Graphs
- Inverse Trigonometric Functions and why . . .

Trigonometric functions can be used to model periodic behavior and applications such as musical notes.




Figure 1.40 The radian measure of angle ACB is the length θ of arc AB on the unit circle centered at C . The value of θ can be found from any other circle, however, as the ratio s/r .




Figure 1.41 An angle θ in standard position.

Graphs of Trigonometric Functions

When we graph trigonometric functions in the coordinate plane, we usually denote the independent variable (radians) by x instead of θ . Figure 1.42 on the next page shows sketches of the six trigonometric functions. It is a good exercise for you to compare these with what you see in a grapher viewing window. (Some graphers have a "trig viewing window.")

EXPLORATION 1 Unwrapping Trigonometric Functions

Set your grapher in **radian mode**, **parametric mode**, and **simultaneous mode** (all three). Enter the parametric equations

$$x_1 = \cos t, \quad y_1 = \sin t \quad \text{and} \quad x_2 = t, \quad y_2 = \sin t.$$

- Graph for $0 \leq t \leq 2\pi$ in the window $[-1.5, 2\pi]$ by $[-2.5, 2.5]$. Describe the two curves. (You may wish to make the viewing window square.)
- Use TRACE to compare the y -values of the two curves.
- Repeat part 2 in the window $[-1.5, 4\pi]$ by $[-5, 5]$, using the parameter interval $0 \leq t \leq 4\pi$.
- Let $y_2 = \cos t$. Use TRACE to compare the x -values of curve 1 (the unit circle) with the y -values of curve 2 using the parameter intervals $[0, 2\pi]$ and $[0, 4\pi]$.
- Set $y_2 = \tan t$, $\csc t$, $\sec t$, and $\cot t$. Graph each in the window $[-1.5, 2\pi]$ by $[-2.5, 2.5]$ using the interval $0 \leq t \leq 2\pi$. How is a y -value of curve 2 related to the corresponding point on curve 1? (Use TRACE to explore the curves.)